

Hypothalamic-immune interactions: neuromodulation of natural killer activity by lesioning of the anterior hypothalamus

R. J. CROSS, W. R. MARKESBERY, W. H. BROOKS & T. L. ROSZMAN *Departments of Medical Microbiology and Immunology, Pathology, Neurology, the Sanders-Brown Research Center on Aging and the Division of Neurosurgery, University of Kentucky Medical Center, Lexington, Kentucky, U.S.A.*

Accepted for publication 22 August 1983

Summary. Placement of bilateral electrolytic lesions in the preoptic-anterior hypothalamic area (AHT) of Fischer 344 rats results in decreased splenic NK activity as compared to control and normal animals. Animals with AHT lesions have a decrease in NK activity 4 and 7 days after lesion placement, with a return to normal activity by day 14. Fractionation of spleen cells on glass bead columns results in non-adherent and adherent cell fractions with NK activity. AHT lesions affect NK activity only in the adherent cell fraction. The removal of macrophages from this cell fraction did not restore NK activity. Moreover, this NK activity is not the result of cytotoxic macrophages. Hypophysectomy decreases NK activity in lesioned and non-lesioned animals, suggesting the influence of pituitary factors. These data indicate that the anterior hypothalamus is capable of modulating the cytotoxic activity of NK cells. Thus, neuroimmunomodulation may be a potential factor in susceptibility to some disease states such as viral infections and neoplasia.

INTRODUCTION

Recently it has been shown that regulation of the immune response is not limited to factors or networks endogenous to the immune system (Benacerraf &

Correspondence: Dr Richard J. Cross, Department of Medical Microbiology and Immunology, University of Kentucky Medical Center, Lexington, KY 40536, U.S.A.

McDevitt, 1972; McDevitt, 1980; Gershon, 1974; Jerne, 1974), but also includes mechanisms that emanate from the central nervous system (CNS) (Besedovsky & Sorkin, 1977). These effectors include hormones (Dougherty *et al.*, 1964; Vischer, 1972; Ambrose, 1970; Filipp & Mess, 1969; Fabris, 1973; Fabris, Pierpaoli & Sorkin, 1971; Pierpaoli & Maestroni, 1977, 1978), neurotransmitters (Devoino *et al.*, 1975; Bliznakov, 1980), endorphins (Gilman *et al.*, 1982; Johnson *et al.*, 1982) or sympathetic nerve pathways (Besedovsky *et al.*, 1979; Kasahara *et al.*, 1977; Williams *et al.*, 1981; Williams & Felten, 1981; Giron, Crutcher & Davis, 1980). Additional support is found for a concept of neural-immune interactions in reports which indicate that perturbations of the CNS by electrolytic lesions in the hypothalamus and the limbic system induce a variety of immune alterations. These include: (i) changes in lymphoid tissue architecture (Isakovic & Jankovic, 1973; Cross *et al.*, 1980; Brooks *et al.*, 1982), (ii) decreased susceptibility to induction of delayed-type hypersensitivity (Jankovic & Isakovic, 1973), (iii) decreased anaphylactic reactions (Luparello, Stein & Park, 1964), (iv) impaired or enhanced non-specific lymphocyte activation (Cross *et al.*, 1980; Brooks *et al.*, 1982; Keller *et al.*, 1980; Cross *et al.*, 1982), (v) decreased responsiveness to transplantation antigens (Dann, Wachtel & Rubin, 1979), and (vi) decreased humoral response to protein antigens (Jankovic & Isakovic, 1973; Tyrey & Nalbandov, 1972).

Several clinical correlates for neuroimmunomodulation also exist. For example, depressed immune responsiveness in patients with cerebral tumours (Brooks *et al.*, 1972), Huntington's disease (Morrell, 1979) or Guamanian-Parkinsonism-dementia complex (Hoffman *et al.*, 1978) indicate that diseases of the CNS are associated with changes in immune competence. Additionally, individuals with psychosis (Kovaleva, Bonartsev & Prilipko, 1977), bereavement and depression (Bartrop, Lazarus & Luckhurst, 1977), and emotional stress also exhibit impaired lymphocyte reactivity and antibody production (Solomon, 1969a). Investigations of experimental animals stressed by overcrowding or avoidance conditioning display impaired immune responsiveness (Rasmussen, Spencer & Marsh, 1959; Solomon, 1969b), decreased resistance to viral infections (Rasmussen, 1969), and an increased incidence of neoplasia (Riley, Fitzmaurice & Spackman, 1981), further suggesting a link between behaviour and disease susceptibility.

Numerous reports suggest that immune surveillance for viral infections and neoplastic disease is provided by either cytotoxic T lymphocytes (Martz, 1974) or natural killer (NK) cells (Herberman & Ortaldo, 1981; Herberman & Holden, 1978). NK cells are cytotoxic for a variety of cells including virally transformed (Williams, Leifer & Moore, 1977), lymphoma (Nunn *et al.*, 1976), and those derived from solid tumours (Brooks *et al.*, 1980).

The purpose of this report is to determine if lesions in the preoptic-anterior hypothalamic area (AHT) modulate NK activity.

MATERIALS AND METHODS

Animals

Hypophysectomized and non-hypophysectomized male Fischer 344 rats (Charles River, Wilmington, MA), age 65–90 days, were used in these experiments. Hypophysectomy (HX) was performed 3–4 weeks prior to use and HX rats were maintained on 5% glucose solution.

Induction of brain lesions

Bilateral, electrolytic lesions were placed in the AHT as reported previously (Cross *et al.*, 1980). Rats with lesions in the frontal grey matter (FC) served as controls. Rats were anaesthetized with chloral hydrate (35 mg/kg). Lesions were placed using direct current (1.5 mA for 15 sec) through a noble-platinum elec-

trode at the following coordinates: AHT—1.2 mm anterior to the bregma, 0.7 mm lateral of the midline and 7.4 mm below the dura; FC—4.0 mm anterior to the bregma, 3.0 mm lateral of the midline and 1.5 mm below the dura.

At the time of death, the brain was removed, fixed in 4% formalin, and sectioned in a coronal plane. Haematoxylin and eosin-stained sections were examined for location and extent of the lesions. All animals whose lesions did not fall within the AHT as previously defined (Cross *et al.*, 1980) were discarded. Mortality was <10% and confined to the first 24 hr following surgery.

Preparation of spleen cell suspensions

Animals were anaesthetized with methoxyflurane (Pitman-Moore Inc., Washington Crossing, NJ) and killed by cervical dislocation. Spleens were removed and teased into dispersed cell suspensions in RPMI-1640 (Gibco, Grand Island, NY). An aliquot was removed and set aside on ice for later evaluation of NK activity. The remainder of the cells were fractionated over a glass bead column, and non-adherent (GBNA) and adherent (GBA) cell fractions were collected (Shortman *et al.*, 1971). Approximately 58% of the cells were recovered in the GBNA cell fraction and 12% were recovered in the GBA cell fraction. An aliquot of adherent cells was incubated with 200 mg of carbonyl iron (Pfaltz and Bauer, Sanford, CT) in 10 ml of RPMI-1640 supplemented with 5% foetal calf serum (FCS; Gibco, Grand Island, NY) for 1 hr at 37°. A magnet was applied to the bottom of the flask and the suspended cells were transferred to plastic 50 ml conical tubes. The iron filings remaining in the flask were resuspended in 10 ml of RPMI-1640 with 5% FCS, the magnet applied, and the cells transferred to a second 50 ml conical tube. The cells were centrifuged, washed, and resuspended in 5 ml of medium. Any remaining iron filings were removed from this cell suspension by employing the magnet, the cells transferred to another tube, washed once and counted. The carbonyl iron-treated cell fraction contained <1% macrophages as determined by non-specific esterase staining (Yam, Li & Crosby, 1971).

Natural killer cell assay

NK activity was quantified using a ⁵¹Cr-release assay. Spleen cells were suspended in RPMI-1640 supplemented with penicillin, streptomycin, vitamins, glutamine, non-essential amino acids, 10% heat-inactivated

FCS and adjusted to 1×10^7 cells/ml. All cell populations were incubated for 2 hr at 37° prior to assay.

Adenovirus type 2 transformed rat fibroblasts (F4) which served as NK target cells (obtained from Dr J. Sheil, University of Kentucky) were labelled with 250 μ Ci sodium 51 chromate for 1 hr at 37° . Target cells were washed three times, counted, and adjusted to 1×10^5 cells/ml. NK activity was measured by adding 100 μ l of F4 target cells to a 96-well round-bottom microtitre plate. Spleen cells and target cells were added at an effector:target ratio of 100:1, 50:1, 25:1, and 12.5:1. Supernatant fluids were harvested after 18 hr using the Titertek supernatant collection system (Flow Laboratories, McLean, VA) and released 51 Cr was counted in a Beckman Gamma 8000. Percent cytotoxicity was calculated using the following formula:

% cytotoxicity =

$$\frac{\text{c.p.m. experimental release} - \text{c.p.m. spontaneous release}}{\text{c.p.m. maximum release} - \text{c.p.m. spontaneous release}} \times 100$$

Lytic units were calculated by fitting regression lines using percent cytotoxicity obtained at each effector:target cell ratio. One lytic unit was arbitrarily set at 20% cytotoxicity and the number of effector cells required to obtain this level of killing was obtained from the regression line. Lytic units/ 10^6 cells were calculated accordingly. Regression lines were generated for all four effector cell populations (unfractionated, GBNA, GBA, and carbonyl iron-treated GBA) from each of the three groups of animals (AHT, FC, and normal). The data were analysed by the two-tailed Student's *t*-test for independent means.

RESULTS

Effects of AHT lesions on NK activity

Experiments were performed to determine the effects of AHT lesions on NK activity. Four days after placement of AHT lesions splenic NK activity was determined. As shown in Fig. 1, rats with AHT lesions display impaired NK activity as compared to FC-lesioned and normal animals. This diminished response persisted through a range of effector:target cell ratios. FC-lesioned animals did not differ from normals in their NK activity.

Kinetics of AHT lesion-induced NK impairment

The kinetics of diminished NK activity were determined in animals killed 4, 7 and 14 days after lesioning.




Figure 1. Splenic NK activity in AHT (Δ), FC (\blacksquare) and normal (\bullet) animals at increasing effector:target cell ratios 4 days after lesion placement. NK activity in AHT-lesioned animals is significantly different from FC-lesioned ($P < 0.01$) and normal animals ($P < 0.001$). NK activity in FC-lesioned animals did not differ from normals.

Previous experiments evaluating the effects of AHT lesions on mitogen-induced lymphocyte proliferation have shown that the maximum effect occurs 4 days after lesioning and returns to normal values by the 14th day (Cross *et al.*, 1980). Figure 2 shows that AHT lesions significantly ($P < 0.01$) depress NK activity at 4 and 7 days. NK activity returns to normal values by the fourteenth day. Although NK activity is enhanced at day 14, it is not significantly different from that seen in normal animals. Hence the kinetics of the impairment of NK activity after AHT lesioning is similar to that observed when assessing the effects of AHT lesions on mitogen-induced lymphocyte proliferation.

Effect of glass bead column fractionation on NK activity

To further evaluate the effects of AHT lesions on NK activity, spleen cells from AHT, FC, or normal animals were fractionated on glass bead columns.




Figure 2. Splenic NK activity in AHT (▲), FC (■) and normal animals 4, 7 and 14 days after lesion placement (values represent 100:1 effector:target cell ratio). NK activity in AHT-lesioned animals is significantly different from FC-lesioned animals ($P < 0.01$) at both days, 4 and 7. NK activity in AHT- or FC-lesioned animals at day 14 did not differ from normal.

After fractionation, GBNA and GBA cell populations were tested for NK activity. NK activity is detected in all cell fractions. As shown in Table 1, AHT lesions impair NK activity in unfractionated spleen cell

populations as compared to FC-lesioned or normal animals. NK activity in the GBNA cell fraction obtained from AHT, FC and normal animals do not differ. However, NK activity in the GBA cell fraction of the AHT-lesioned group is decreased when compared to that obtained from either FC-lesioned or normal animals. These data indicate that AHT lesions affect NK activity only in the GBA cell population.

Recent studies using a long-term 18 hr ^{51}Cr -release assay have shown that natural cytotoxic activity in rats may be due to cytotoxic macrophage activity (Brooks *et al.*, 1980). To investigate this, GBA spleen cells from AHT, FC, and normal animals were treated with carbonyl iron to remove macrophages. This treatment decreases the number of non-specific esterase positive cells from 12% to <1%. Removal of these macrophages (Table 1) did not alter the level of cytotoxicity within the GBA cell fraction of each group, indicating that cytotoxic macrophages are not involved in the lysis of F4 target cells. We have previously shown that AHT lesions enhance suppressor macrophage activity which is capable of inhibiting mitogen-induced lymphocyte proliferation (Roszman *et al.*, 1982). NK activity in AHT-lesioned animals continues to be impaired despite the removal of macrophages by carbonyl iron, thus, indicating that diminished NK activity does not result from enhanced macrophage suppression.

NK activity in hypophysectomized rats with AHT lesions

The next set of experiments were performed to determine the role of the pituitary gland in the

Table 1. NK activity of non-fractionated and glass bead column fractionated spleen cells from normal and FC-, AHT-lesioned animals*

Groups	Lytic units/ 10^6 cells \pm SEM			
	Unfractionated	GBNA	GBA	Carbonyl iron-treated GBA
AHT Lesions ($n = 16$) [‡]	1.46 ± 0.06 ($P < 0.01$) [†]	1.59 ± 0.16 ($P > 0.05$)	1.42 ± 0.19 ($P < 0.05$)	1.45 ± 0.32 ($P < 0.05$)
FC Lesions ($n = 18$)	2.62 ± 0.62	1.61 ± 0.11	2.25 ± 0.25	2.75 ± 0.24
Normal ($n = 18$)	2.50 ± 0.30	1.71 ± 0.15	3.42 ± 0.67	2.86 ± 0.33

* Spleen cells obtained from animals 4 days after AHT or FC lesion placement.

[†] Significance was determined by Student's *t*-test comparing values obtained from lesioned animals to those from control animals. Normal and FC-lesioned animals did not differ significantly.

[‡] n = number of animals/group.

Table 2. The effect of AHT lesions on NK activity in hypophysectomized animals*

Group	Lytic units/10 ⁶ cells \pm SEM	
	Non-hypophysectomized†	Hypophysectomized‡
AHT	1.46 \pm 0.06 (n=16)‡	1.01 \pm 0.06§ (n=8)
FC	2.62 \pm 0.62 (n=18)	1.24 \pm 0.08 (n=10)
Normal	2.50 \pm 0.30 (n=18)	1.38 \pm 0.13 (n=6)

* NK activity determined 4 days after lesioning.

† NK activity in the hypophysectomized groups differs significantly from that of non-hypophysectomized groups ($P < 0.02$).

‡ n = number of animals/group.

§ NK activity did not differ significantly among the hypophysectomized groups.

suppression of NK activity after AHT lesioning. NK activity in all HX groups is significantly less than in animals with an intact pituitary (Table 2). These data indicate that hypophysectomy has a marked effect in impairing NK activity as previously reported (Saxena, Saxena & Adler, 1982). Furthermore, these observations suggest that pituitary factors are important for the induction of NK impairment by AHT lesions.

DISCUSSION

The most meaningful finding of this study is that electrolytic lesioning of the anterior hypothalamic area results in depressed NK activity. This is the first indication that discrete lesions in the hypothalamus affect the NK cell which may be important in immune surveillance. These data correlate with our previous reports on the effects of AHT lesions on mitogen-induced lymphocyte proliferation (Cross *et al.*, 1980; Brooks *et al.*, 1982). It is unlikely that the effect of AHT lesions on NK activity is the result of increased circulating corticosteroid levels because we have observed that both AHT and FC-lesioned animals have similar levels (Cross *et al.*, 1982). The transitory nature of these effects may result from either neuronal plasticity (Goldowitz, Scheff & Cotman, 1979) or the rapid turn-over of neurally controlled factors (e.g. hormones, peptides) which may serve as mediators of CNS modulation of immunity.

Recently Renoux *et al.* (1982) have shown that unilateral removal of the left neocortex impairs NK activity. Although small, discrete bilateral electrolytic

lesions placed in the frontal cortex are associated with a slight decrease in NK activity, the suppression is not significant when compared to normal animals. These differences may result from the size and extent of the lesions or the method in which the lesions were produced (surgical *vs* electrolytic).

Previously we have demonstrated that AHT lesions enhance macrophage suppressor cell function (Roszman *et al.*, 1982). Hochman, Cudkowicz & Evans (1981) have shown that removal of macrophages by carbonyl iron increases splenic NK activity. The present report indicates that removal of macrophages by carbonyl iron does not alleviate the diminished NK activity resulting from AHT lesioning. Further experiments demonstrated that spleen cells can be fractionated to yield non-adherent and adherent cell fractions with NK activity. Interestingly, hypothalamic lesions employed in this study affect NK activity only in the adherent cell fraction, whereas activity in the non-adherent fraction is not altered. The diminished NK activity in the adherent fraction is not restored after removal of the macrophage. These findings indicate that macrophage suppression is not responsible for altered NK function. Moreover, in the present study, killing by the adherent population is not the result of macrophage-mediated cell lysis.

An intact pituitary is necessary for the maintenance of NK activity. Saxena, Saxena & Adler (1982) have shown that NK function is depressed in mice after hypophysectomy. Reconstitution with growth hormone restores NK activity to normal levels in these animals. Alternatively, Hanna & Schneider (1983) have shown that the steroid hormone, β -estradiol, suppresses NK activity. Recently, it has been shown that β -endorphins enhance NK activity in humans (J. E. Morley, personal communication). These studies lend support to the theory that hormones and neuro-peptides modulate the NK cell. The present study confirms that hypophysectomy markedly decreases NK activity. Decreased NK activity in AHT lesioned animals is abrogated by hypophysectomy, suggesting that neural modulation of NK cells is affected by pituitary hormones or peptides. This conclusion must be tempered, however, by the observation that hypophysectomy has such a profound effect on NK activity that further modulation of NK function by non-pituitary factors (e.g. neurotransmitters) would be difficult to detect in this system. Further studies are needed to delineate the role of hormones, neurotransmitters and neuropeptides as modulators of NK cells.

In summary, the present study indicates that the

CNS, particularly the anterior hypothalamus, is capable of modulating the cytolytic activity of adherent NK cells. Thus, neuroimmunomodulation may be a potential factor in susceptibility to certain diseases.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the expert technical assistance of Donna Alexander. This work was supported in part by N.I.H. grants CA-18234, NS-17423 and AG 02435.

REFERENCES

- AMBROSE C.T. (1970) Hormones and the immune response. In: *Ciba Foundation Study Group No. 36*, p. 100. J. and A. Churchill, London.
- BARTROP R.W., LAZARUS E. & LUCKHURST L.G. (1977) Depressed lymphocyte function after bereavement. *Lancet*, **i**, 834-836.
- BENACERRAF B. & McDEVITT H.O. (1972) Histocompatibility-linked immune response genes. *Science*, **175**, 273.
- BESEDOVSKY H.O., DEL RAY A., SORKIN E., DA PRADA M. & KELLER H.H. (1979) Immunoregulation mediated by the sympathetic nervous system. *Cell. Immunol.* **48**, 346.
- BESEDOVSKY H.O. & SORKIN E. (1977) Network of immune-neuroendocrine interactions. *Clin. exp. Immunol.* **27**, 1.
- BLIZNAKOV E. (1980) Serotonin and its precursors as modulators of immunological responsiveness in mice. *J. Med.* **11**, 81.
- BROOKS C.G., FLANNER G.R., WEBB P.J. & BALDWIN R.W. (1980) Quantitative studies of natural cytotoxicity to solid tumors in rats. The nature of the killer cell depends on the type of assay. *Immunology*, **41**, 673.
- BROOKS W.H., CROSS R.J., ROSZMAN T.L. & MARKESBERY W.R. (1982) Neuroimmunomodulation: neural anatomical basis for impairment and facilitation. *Ann. Neurol.* **12**, 56.
- BROOKS W.H., NETSKY M.G., NORMANSELL D.E. & HOROWITZ D.A. (1972) Depressed cell-mediated immunity in patients with primary intracranial tumors. *J. exp. Med.* **136**, 1631.
- CROSS R.J., BROOKS W.H., ROSZMAN T.L. & MARKESBERY W.R. (1982) Hypothalamic-immune interactions. Effect of hypophysectomy on neuroimmunomodulation. *J. Neurol. Sci.* **53**, 557.
- CROSS R.J., MARKESBERY W.R., BROOKS W.H. & ROSZMAN T.L. (1980) Hypothalamic-immune interactions. I. The acute effect of anterior hypothalamic lesions on the immune response. *Brain Res.* **196**, 79.
- DANN J.A., WACHTEL S.S. & RUBIN A.L. (1979) Possible involvement of the central nervous system in graft rejection. *Transplant.* **27**, 223.
- DEVOINO L., ELISEVA L., EREMINA O., IDOVA G. & CHEIDO M. (1975) 5-Hydroxytryptophan effect on the development of the immune response: IgM and IgG antibodies and rosette formation in primary and secondary responses. *Eur. J. Immunol.* **5**, 394.
- DOUGHERTY T.F., BERLINER M.L., SCHNEEBELI G.L. & BERLINER D.L. (1964) Hormonal control of lymphatic structure and function. *Ann. N.Y. Acad. Sci.* **113**, 825.
- FABRIS N. (1973) Immunodepression in thyroid-deprived animals. *Clin. exp. Immunol.* **15**, 601.
- FABRIS N., PIERPAOLI W. & SORKIN E. (1971) Hormones and the immunological capacity. IV. Restorative effects of developmental hormones or of lymphocytes on the immunodeficiency syndrome of the dwarf mouse. *Clin. exp. Immunol.* **9**, 227.
- FILIPP G. & MESS B. (1969) Role of the thyroid hormone system in suppression of anaphylaxis due to electrolytic lesion of the tuberal region of the hypothalamus. *Ann. Allergy*, **27**, 500.
- GERSHON R.K. (1974) T-cell control of antibody responses. *Contemp. Top. Immunobiol.* **3**, 1.
- GILMAN S.C., SCHWARTZ J.M., MILNER R.J., BLOOM F.E. & FELDMAN J.D. (1982) β -Endorphin enhances lymphocyte proliferative responses. *Proc. natn. Acad. Sci. U.S.A.* **79**, 4226.
- GIRON L.T., CRUTCHER K.A. & DAVIS J.N. (1980) Lymph nodes—a possible site for sympathetic neuronal regulation of immune responses. *Ann. Neurol.* **8**, 520.
- GOLDOWITZ D., SCHEFF S. & COTMAN C. (1979) The specificity of reactive synaptogenesis: comparative study in the adult rat hippocampal formation. *Brain Res.* **170**, 427.
- HANNA N. & SCHNEIDER M. (1983) Enhancement of tumor metastasis and suppression of natural killer cell activity by β -estradiol treatment. *J. Immunol.* **130**, 974.
- HERBERMAN R.B. & HOLDEN H.T. (1978) Natural cell-mediated immunity. *Adv. Cancer Res.* **27**, 305.
- HERBERMAN R.B. & ORTALDO J.R. (1981) Natural killer cells: their role in defense against disease. *Science*, **214**, 24.
- HOCHMAN P.S., CUDKOWICZ G. & EVANS P.D. (1981) Carageenan-induced decline of natural killer activity. II. Inhibition of cytotoxicity by adherent non-T 1a-negative suppressor cells activated *in vivo*. *Cell. Immunol.* **61**, 200.
- HOFFMAN P.M., ROBBINS D.S., NOLTE M.T., GIBBS C.J. & GAJDUSEK D.C. (1978) Immunity and immunogenetics in Guamanians with amyotrophic lateral sclerosis and Parkinsonism-dementia complex. *N. Engl. J. Med.* **299**, 680.
- ISAKOVIC K. & JANKOVIC B.D. (1973) Neuroendocrine correlates of immune response. II. Changes in the lymphatic organs of brain-lesioned rats. *Int. Archs Allergy appl. Immun.* **45**, 373.
- JANKOVIC B.D. & ISAKOVIC K. (1973) Neuroendocrine correlates of immune response. I. Effects of brain lesions on antibody production, Arthus reactivity, and delayed hypersensitivity in the rat. *Int. Archs Allergy appl. Immun.* **45**, 360.
- JERNE N.K. (1974) Towards a theory of the immune system. *Ann. Immunol. (Inst. Pasteur)*, **125C**, 373.
- JOHNSON H.M., SMITH E.M., TORRES B.A. & BLALOCK J.E. (1982) Regulation of the *in vitro* antibody response by neuroendocrine hormones. *Proc. natn. Acad. Sci. U.S.A.* **79**, 4171.
- KASAHARA K., TANAKA S., ITO T. & HAMASHIMA Y. (1977) Suppression of the primary immune response by chemical sympathectomy. *Res. Commun. Chem. Pathol. Pharmacol.* **16**, 687.

- KELLER S.E., STEIN M., CAMERINO M.S., SCHLEIFER M.S. & SHERMAN J. (1980) Suppression of lymphocyte stimulation by anterior hypothalamic lesions in the guinea pig. *Cell. Immunol.* **52**, 557.
- KOVALEVA E.S., BONARTSEV P.D. & PRILIPKO L.L. (1977) Lymphocyte reaction in schizophrenic patients to the phytomitogens, concanavalin A and phytohemagglutinin. *Bull. exp. Biol. Med.* **84**, 1136.
- LUPARELLO T.J., STEIN M. & PARK C.D. (1964) Effect of hypothalamic lesions on rat anaphylaxis. *Am. J. Physiol.* **207**, 911.
- MARTZ E. (1974) Mechanism of specific tumor-cell lysis by alloimmune T-lymphocytes: resolution and characterization by discrete steps in the cellular interaction. *Contemp. Top. Immunobiol.* **7**, 301.
- MCDEVITT H.O. (1980) Regulation of the immune response by the major histocompatibility system. *N. Engl. J. Med.* **303**, 1514.
- MORRELL R.M. (1979) Antibody-dependent cytotoxicity in Huntington's disease. *Adv. Neurol.* **23**, 443.
- NUNN M.E., DJEU J.Y., GLASER M., LAVRIN P.H. & HERBERMAN R.B. (1976) Natural cytotoxic reactivity of rat lymphocytes against syngeneic Gross virus-induced lymphoma. *J. natn. Cancer. Inst.* **56**, 393.
- PIERPAOLI W. & MAESTRONI G.J.M. (1977) Pharmacological control of the immune response by blockade of the early hormonal changes following antigen injections. *Cell. Immunol.* **31**, 355.
- PIERPAOLI W. & MAESTRONI G.J.M. (1978) Pharmacologic control of the hormonally-modulated immune response. II. Blockade of antibody production by a combination of drugs acting on neuroendocrine functions: its prevention by gonadotropins and corticotropin. *Immunology*, **34**, 419.
- RASSMUSSEN A.F. (1969) Emotions and immunity. *Ann. N.Y. Acad. Sci.* **164**, 458.
- RASSMUSSEN A.F., SPENCER E.S. & MARSH J.T. (1959) Decreased susceptibility of mice to passive anaphylaxis following avoidance-learning stress. *Proc. Soc. exp. Biol. Med.* **100**, 878.
- RENOUX G., BIZIERE K., BARDOS P., DEGENNE D. & RENOUX M. (1982) NK activity in mice is controlled by the brain neocortex. IN: *NK Cells and Other Effector Cells*, p. 639. Academic Press, New York.
- RILEY V., FITZMAURICE M.A. & SPACKMAN D. (1981) Psychoneuroimmunologic factors in neoplasia: studies in animals. In: *Psychoneuroimmunology*, p. 31. Academic Press, New York.
- ROSZMAN T.L., CROSS R.J., BROOKS W.H. & MARKESBERY W.R. (1982) Hypothalamic-immune interactions. II. The effect of hypothalamic lesions on the ability of adherent spleen cells to limit lymphocyte blastogenesis. *Immunology*, **45**, 737.
- SAXENA Q.B., SAXENA R.K. & ADLER W.H. (1982) Regulation of natural killer cell activity *in vivo*. III. Effect of hypophysectomy and growth hormone treatment on the natural killer activity of the mouse spleen cell population. *Int. Archs Allergy appl. Immun.* **67**, 169.
- SHORTMAN K., WILLIAMS N., JACKSON H., RUSSELL P., BYRT P. & DIERNER E. (1971) The separation of different cell classes from lymphoid organs. IV. The separation of lymphocytes from phagocytes on glass bead columns, and its effect on subpopulations of lymphocytes and antibody-forming cells. *J. Cell. Biol.* **48**, 566.
- SOLOMON G.F. (1969a) Emotions, stress, the central nervous system and immunity. *Ann. N.Y. Acad. Sci.* **164**, 335.
- SOLOMON G.F. (1969b) Stress and antibody response in rats. *Int. Archs Allergy appl. Immun.* **35**, 97.
- TYREY L. & NALBANDOV V.P. (1972) Influence of anterior hypothalamic lesions on circulating antibody titers in the rat. *Am. J. Physiol.* **222**, 179.
- VISCHER T.L. (1972) Effect of hydrocortisone on the reactivity of thymus and spleen cells of mice to *in vitro* stimulation. *Immunology*, **23**, 777.
- WILLIAMS J.M. & FELTEN D.L. (1981) Sympathetic innervation of murine thymus and spleen: a comparative histofluorescence study. *Anatom. Rec.* **199**, 531.
- WILLIAMS J.M. & PETERSON R.G., SHEA P.A., SCHMEDITJE J.F., BAUER D.C. & FELTEN D.L. (1981) Sympathetic innervation of murine thymus and spleen: evidence for a functional link between the nervous and immune systems. *Brain Res. Bull.* **6**, 83.
- WILLIAMS R.M., LEIFER J. & MOORE M.J. (1977) Hybrid effect in natural cell-mediated cytotoxicity of SV-40 transformed fibroblasts by rat spleen cells. *Transplant.* **23**, 283.
- YAM L.T., LI C.Y. & CROSBY W.H. (1971) Cytochemical identification of monocytes and granulocytes. *Amer. J. clin. Pathol.* **55**, 283.